


Claims

1. A method of treatment of bacterial infections in mammals, which method comprises the administration to a mammal in need of such treatment of an effective amount of a compound of formula (I) or a pharmaceutically acceptable derivative thereof:



(I)

wherein:

one of Z¹, Z², Z³, Z⁴ and Z⁵ is N or CR^{1a} and the remainder are CH;

R¹ is selected from hydroxy; (C₁₋₆) alkoxy optionally substituted by (C₁₋₆)alkoxy, amino, piperidyl, guanidino or amidino optionally N-substituted by one or two (C₁₋₆)alkyl, acyl or (C₁₋₆)alkylsulphonyl groups, NH₂CO, hydroxy, thiol, (C₁₋₆)alkylthio, heterocyclylthio, heterocyclyloxy, arylthio, aryloxy, acylthio, acyloxy or (C₁₋₆)alkylsulphonyloxy; (C₁₋₆)alkoxy-substituted (C₁₋₆)alkyl; halogen; (C₁₋₆)alkyl; (C₁₋₆)alkylthio; trifluoromethyl; nitro; azido; acyl; acyloxy; acylthio; (C₁₋₆)alkylsulphonyl; (C₁₋₆)alkylsulphoxide; arylsulphonyl; arylsulphoxide or an amino, piperidyl, guanidino or amidino group optionally N-substituted by one or two (C₁₋₆)alkyl, acyl or (C₁₋₆)alkylsulphonyl groups, or when one of Z¹, Z², Z³, Z⁴ and Z⁵ is N, R¹ may instead be hydrogen;

R^{1a} is selected from hydrogen and the groups listed above for R¹;

R³ is in the 2- or 3-position and is:

carboxy; (C₁₋₆)alkoxycarbonyl; aminocarbonyl wherein the amino group is optionally substituted by hydroxy, (C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl, aminocarbonyl(C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkylsulphonyl, trifluoromethylsulphonyl, (C₁₋₆)alkenylsulphonyl, (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl or (C₂₋₆)alkenylcarbonyl and optionally further substituted by (C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl, aminocarbonyl(C₁₋₆)alkyl or (C₂₋₆)alkenyl; cyano; tetrazolyl; 2-oxo-oxazolidinyl optionally substituted by R¹⁰; 3-hydroxy-3-cyclobutene-1,2-dione-4-yl; 2,4-

thiazolidinedione-5-yl; tetrazol-5-ylaminocarbonyl; 1,2,4-triazol-5-yl optionally substituted by R¹⁰; or 5-oxo-1,2,4-oxadiazol-3-yl; or

R³ is in the 2- or 3-position and is (C₁₋₄)alkyl or ethenyl substituted with any of the groups listed above for R³ and 0 to 2 groups R¹² independently selected from:

- 5 thiol; halogen; (C₁₋₆)alkylthio; trifluoromethyl; azido; (C₁₋₆)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; (C₂₋₆)alkenyloxycarbonyl; (C₂₋₆)alkenylcarbonyl; hydroxy optionally substituted by (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl, (C₂₋₆)alkenylcarbonyl or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkylcarbonyl or (C₂₋₆)alkenylcarbonyl; amino optionally mono- or disubstituted by
- 10 (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl, (C₂₋₆)alkenylcarbonyl, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkylsulphonyl, (C₂₋₆)alkenylsulphonyl or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl; aminocarbonyl wherein the amino group is optionally
- 15 substituted by (C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl, aminocarbonyl(C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl or (C₂₋₆)alkenylcarbonyl and optionally further substituted by (C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl, aminocarbonyl(C₁₋₆)alkyl or (C₂₋₆)alkenyl; oxo; (C₁₋₆)alkylsulphonyl; (C₂₋₆)alkenylsulphonyl; or (C₁₋₆)aminosulphonyl wherein the amino group is optionally
- 20 substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl;
provided that when R³ is disubstituted with hydroxy or amino and carboxy containing substituents these may optionally together form a cyclic ester or amide linkage, respectively;
and provided that R³ is other than (C₁₋₄)alkyl or ethenyl substituted by (C₁₋₆)alkoxycarbonyl or aminocarbonyl optionally substituted by (C₁₋₆)alkyl, (C₂₋₆)alkenyl,
- 25 (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl or (C₂₋₆)alkenylcarbonyl and optionally further substituted by (C₁₋₆)alkyl, hydroxy(C₁₋₆)alkyl, aminocarbonyl(C₁₋₆)alkyl or (C₂₋₆)alkenyl and 0 to 2 groups R¹²;
- 30 wherein R¹⁰ is selected from (C₁₋₄)alkyl; (C₂₋₄)alkenyl; aryl; a group R¹² as defined above; carboxy; aminocarbonyl wherein the amino group is optionally substituted by hydroxy, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkylsulphonyl, trifluoromethylsulphonyl, (C₁₋₆)alkenylsulphonyl, (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl or (C₂₋₆)alkenylcarbonyl and optionally further substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl; cyano; or tetrazolyl;
- 35 (C₁₋₆)alkyl or (C₂₋₆)alkenyl; cyano; or tetrazolyl;

R⁴ is a group -CH₂-R⁵ in which R⁵ is selected from:

(C₃₋₁₂)alkyl; hydroxy(C₃₋₁₂)alkyl; (C₁₋₁₂)alkoxy(C₃₋₁₂)alkyl; (C₁₋₁₂)alkanoyloxy(C₃₋₁₂)alkyl; (C₃₋₆)cycloalkyl(C₃₋₁₂)alkyl; hydroxy-, (C₁₋₁₂)alkoxy- or (C₁₋₁₂)alkanoyloxy-(C₃₋₆)cycloalkyl(C₃₋₁₂)alkyl; cyano(C₃₋₁₂)alkyl; (C₂₋₁₂)alkenyl; (C₂₋₁₂)alkynyl; tetrahydrofuryl; mono- or di-(C₁₋₁₂)alkylamino(C₃₋₁₂)alkyl; acylamino(C₃₋₁₂)alkyl; (C₁₋₁₂)alkyl- or acyl-aminocarbonyl(C₃₋₁₂)alkyl; mono- or di-(C₁₋₁₂)alkylamino(hydroxy) (C₃₋₁₂)alkyl; optionally substituted phenyl(C₁₋₂)alkyl, phenoxy(C₁₋₂)alkyl or phenyl(hydroxy)(C₁₋₂)alkyl; optionally substituted diphenyl(C₁₋₂)alkyl; optionally substituted phenyl(C₂₋₃)alkenyl; optionally substituted benzoyl or benzoylmethyl; optionally substituted heteroaryl(C₁₋₂)alkyl; and optionally substituted heteroaroyl or heteroaroylmethyl;

n is 0, 1 or 2;

either A-B is NHC(O)NH or NHC(O)O, or

A is NR¹¹, O, S(O)_x or CR⁶R⁷ and B is NR¹¹, O, S(O)_x or CR⁸R⁹ where x is 0, 1 or 2 and wherein:

each of R⁶ and R⁷ R⁸ and R⁹ is independently selected from: H; thiol; (C₁₋₆)alkylthio; halo; trifluoromethyl; azido; (C₁₋₆)alkyl; (C₂₋₆)alkenyl; (C₁₋₆)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; (C₂₋₆)alkenyloxycarbonyl; (C₂₋₆)alkenylcarbonyl; hydroxy, amino or aminocarbonyl optionally substituted as for corresponding substituents in R³; (C₁₋₆)alkylsulphonyl; (C₂₋₆)alkenylsulphonyl; or (C₁₋₆)aminosulphonyl wherein the amino group is optionally substituted by (C₁₋₆)alkyl or (C₁₋₆)alkenyl; or R⁶ and R⁸ together represent a bond and R⁷ and R⁹ are as above defined; or R⁶ and R⁸ together represent -O- and R⁷ and R⁹ are both hydrogen; or R⁶ and R⁷ or R⁸ and R⁹ together represent oxo; and each R¹¹ is independently H, trifluoromethyl, (C₁₋₆)alkyl, (C₁₋₆)alkenyl, (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₁₋₆)alkenyloxycarbonyl, (C₂₋₆)alkenylcarbonyl, (C₁₋₆)alkyl or (C₁₋₆)alkenyl and optionally further substituted by (C₁₋₆)alkyl or (C₁₋₆)alkenyl;

provided that A and B cannot both be selected from NR¹¹, O and S(O)_x and when one of A and B is CO the other is not CO, O or S(O)_x.

2. A compound of formula (IA) or a pharmaceutically acceptable derivative thereof which is a compound of formula (I) as defined in claim 1 wherein R^3 is other than (C_{1-6}) alkoxycarbonyl; optionally substituted aminocarbonyl, CN or COOH.
3. A compound according to claim 2 wherein Z^5 is CH or N and Z^1 - Z^4 are each CH.
4. A compound according to claim 2 or 3 wherein R^1 is methoxy, amino- or guanidino- (C_{3-5}) alkyloxy, guanidino (C_{3-5}) alkyloxy, piperidyl (C_{3-5}) alkyloxy, nitro or fluoro, and R^{1a} is hydrogen.
5. A compound according to any of claims 2 to 4 wherein R^3 is in the 3-position and is CH_2CO_2H or 2-oxo-oxazolidinyl.
6. A compound according to any of claims 2 to 5 wherein $AB(CH_2)_n$ is $(CH_2)_3$.
7. A compound according to any of claims 2 to 6 wherein R^4 is (C_{5-10}) alkyl, unsubstituted phenyl (C_{2-3}) alkyl or unsubstituted phenyl (C_{3-4}) alkenyl.
8. A compound of formula (I) as defined in claim 1 selected from:
- [3R, 4R]-1-Heptyl-3-(1-(R or S)-hydroxy-2-cyanoethyl)-4-[3-(6-methoxyquinolin-4-yl)propyl]piperidine;
- [3R, 4R]-1-Heptyl-3-(2-(R or S)-oxo-oxazolidin-5-yl)-4-[3-(6-methoxyquinolin-4-yl)propyl]piperidine;
- [3R, 4R]-1-Heptyl-3-(2-cyanoethyl)-4-[3-(6-methoxyquinolin-4-yl)propyl]piperidine;
- [3R, 4R]-1-Heptyl-3-(3-carboxyethyl)-4-[3-(6-methoxyquinolin-4-yl)propyl]piperidine;
- [3R, 4R]-1-Heptyl-3-carboxy-4-[3-(6-methoxyquinolin-4-yl)propyl]piperidine;
- [3R, 4R]-1-Heptyl-3-(carboxymethyl)-4-[3-(6-methoxyquinolin-4-yl)propyl]piperidine;
- [3R, 4R]-1-Heptyl-3-(1-(R or S)-hydroxy-2-carboxyethyl)-4-[3-(6-methoxyquinolin-4-yl)propyl]piperidine;
- [3R, 4R]-1-Heptyl-3-(2-(E)-carboxyethenyl)-4-[3-(6-methoxyquinolin-4-yl)propyl]piperidine;
- N-(cis-3-(R/S)-Ethoxycarbonyl-1-heptyl-4-(S/R)-piperidyl)-N'-(6-methoxyquinolin-4-yl)urea;
- N-(cis-3-(R/S)-Ethoxycarbonyl-1-heptyl-4-(S/R)-piperidyl)-N'-(6-methoxy-[1,5]-naphthyridin-4-yl)urea;
- N-(cis-3-(R/S)-Aminocarbonyl-1-heptyl-4-(S/R)-piperidyl)-N'-(6-methoxy-[1,5]-naphthyridin-4-yl)urea;
- [3R, 4R]-1-Heptyl-4-[3-(R/S)-hydroxy-3-(6-methoxyquinolin-4-yl)propyl]-3-(2-(R or S)-oxo-oxazolidin-5-yl)-piperidine;
- [3R, 4R]-1-Heptyl-3-cyanomethyl-4-[3-(R/S)-hydroxy-3-(6-methoxyquinolin-4-yl)propyl]piperidine;

[3R, 4R]-1-Heptyl-3-cyanomethyl-4-(2-(R)-hydroxy-3-(6-methoxyquinolin-4-yl)propyl)piperidine;

N-(cis-3-(R/S)-Carboxy-1-heptyl-4-(S/R)-piperidyl)-N'-(6-methoxyquinolin-4-yl)urea;

cis-3-(R/S)-Ethoxycarbonyl-1-heptyl-4-(S/R)-(6-methoxyquinolin-4-yl)aminocarbonyloxypiperidine;


cis-3-(R/S)-Carboxy-1-heptyl-4-(S/R)-(6-methoxyquinolin-4-yl)aminocarbonyloxypiperidine;

a compound 18-36 from Table 1;

or a pharmaceutically acceptable derivative of any of the foregoing compounds.

9. A process for preparing compounds of formula (IA) as defined in claim 2, or a pharmaceutically acceptable derivative thereof, which process comprises:

(a) reacting a compound of formula (IV) with a compound of formula (V):




wherein Z^1 , Z^2 , Z^3 , Z^4 and Z^5 , m , n , R^1 , R^2 , R^3 and R^4 are as defined in formula (I), and X and Y may be the following combinations:

- (i) X is M and Y is $CH_2CO_2R^X$
- (ii) X is CO_2RY and Y is $CH_2CO_2R^X$
- (iii) one of X and Y is $CH=SPh_2$ and the other is CHO
- (iv) X is CH_3 and Y is CHO
- (v) X is CH_3 and Y is CO_2R^X
- (vi) X is CH_2CO_2RY and Y is CO_2R^X
- (vii) X is $CH=PR^Z_3$ and Y is CHO
- (viii) X is CHO and Y is $CH=PR^Z_3$
- (ix) X is halogen and Y is $CH=CH_2$
- (x) one of X and Y is COW and the other is $NHR^{11'}$ or NCO
- (xi) one of X and Y is $(CH_2)_p-V$ and the other is $(CH_2)_qNHR^{11'}$, $(CH_2)_qOH$, $(CH_2)_qSH$ or $(CH_2)_qSCOR^X$ where $p+q=1$
- (xii) one of X and Y is CHO and the other is $NHR^{11'}$
- (xiii) one of X and Y is OH and the other is $-CH=N_2$


in which V and W are leaving groups, R^X and R^Y are (C_{1-6}) alkyl and R^Z is aryl or (C_{1-6}) alkyl, or

(xiv) X is NCO, Y is OH or NH_2 ;

- 5 (b) reacting a compound of formula (IV) with a compound of formula (Vb):




(IV)



(Vb)

- 10 wherein Z^1, Z^2, Z^3, Z^4 and Z^5 , m, n, R^1, R^2, R^3 and R^4 are as defined in formula (I), X is $CH_2NHR^{11'}$ and Y is CHO or COW or X is CH_2OH and Y is $-CH=N_2$;


(c) rearranging a compound of formula (II):



(II)

- 15 to give a compound of formula (III) which is a compound of formula (I) where Z^1-Z^5 are CH, n is 1, A-B is $COCH_2$ and R^2 is H, or a compound of formula (VII) which is a compound of formula (I) where n is 1, A-B is $CHOHCH_2$ or CH_2CHOH and R^2 is H; or

(d) photooxygenating a compound of formula (VI):



(VI)

20 in which $Z^{1'}-Z^{5'}$ are Z^1-Z^5 or groups convertible thereto, $R^{11'}, R^1, R^{2'}, R^{3'}$ and $R^{4'}$ are R^{11}, R^1, R^2, R^3 and R^4 or groups convertible thereto, and thereafter optionally or as

necessary converting $R^{11'}$, $R^{1'}$, $R^{2'}$, $R^{3'}$ and $R^{4'}$ to R^{11} , R^1 , R^2 , R^3 and R^4 , converting $Z^{1'}$ - $Z^{5'}$ to Z^1 - Z^5 , converting A-B to other A-B, interconverting R^{11} , R^1 , R^2 , R^3 and/or R^4 and forming a pharmaceutically acceptable derivative thereof.

5 10. A pharmaceutical composition comprising a compound of formula (IA) as defined in claim 2, or a pharmaceutically acceptable derivative thereof, and a pharmaceutically acceptable carrier.

11. The use of a compound of formula (I) as defined in claim 1 or a pharmaceutically acceptable derivative thereof in the manufacture of a medicament for use in the treatment of bacterial infections in mammals.

10

12. A pharmaceutical composition for use in the treatment of bacterial infections in mammals comprising a compound of formula (I) as defined in claim 1, or a pharmaceutically acceptable derivative thereof, and a pharmaceutically acceptable carrier.

ADD
A3

0989820-092001

ADD
A1